



Plagiarism Checker X Originality Report

Similarity Found: 12%

Date: Senin, Agustus 06, 2018

Statistics: 333 words Plagiarized / 2694 Total words

Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

THE EFFECT OF FISHMEAL ON THE LEVELS OF ZINC, VITAMIN A SERUM, NUTRITIONAL INTAKE AND SHORTNESS STATUS OF CHILDREN AGED 2-4 YEARS AFTER ADMINISTRATION OF VITAMIN A IN MAKASSAR, SOUTH SULAWESI, INDONESIA Lydia Fanny¹ Department of Nutrition, Makassar Health Polytechnic, South Sulawesi; Email Address: lydiafanny90@yahoo.com ABSTRACT_ One kind of individual nutrition services offered is supplementation of zinc (Zn) which is recommended to overcome the problem of growth among children.

Animal foods which are source of zinc, such as fish, has a high quality of protein, with digestibility as of 96%. High digestibility of protein is necessary for growth and intelligence. This study aim to the effect of fishmeal on the levels of zinc, vitamin A serum, nutritional intake and shortness status of children aged 2-4 years after administration of vitamin A. This study used experimental study design using Pre-Test and Post-Test Control Group Design.

Sample size was 30 individuals, randomly selected among children aged 2-4 years with shortness status after receiving vitamin A supplementation, which was then divided into two groups of treatments: the group of vitamin A, and the group of vitamin A plus fishmeal. Nutritional status was known through anthropometric measurements using the seca and microtoise for weight and height respectively.

The tools used were in state of good for use and was already been calibrated. Nutritional intake was obtained using a 24-hour recall with consumer multi-pass method. Enumerator were nutritionists, the supervisor was a professor of nutrition at Health Polytechnic Makassar. The study used an internal ethics committee approval at the department of nutrition. Statistical analysis was done using paired t-test with 95%

significance.

Nutritional intake in group I (Vitamin A), all increased after the intervention except for energy intake. Significance values for energy, protein, fat, carbohydrate, Vitamin A and Zn were (0.278) (0.000) (0.000) (0.010) (0.000) and (0.002) respectively. Nutritional intake in group II (Vitamins A, plus fishmeal), all increased after intervention except energy intake.

Significance values for energy, protein, fat, carbohydrate, Vitamin A and Zn were (0.100) (0.000) (0.000) (0.000) (0.000) and (0.001) respectively. Zn levels after intervention in both groups increased significantly. The only obvious difference was in increased levels of albumin which occurred in group II (Vitamin A Plus fishmeal) ($p = 0.010$).

There was no increase in total protein levels in group II (Vitamin A plus fishmeal, while in group I, it was found a significant elevated level of total protein ($p = 0.007$). There was an increase in hemoglobin levels in group II but not in group I. Recommendation from the study: it is necessary to perform nutritional services efforts by providing zinc supplementation in children that suffered shortness status after receiving vitamin A.

INTRODUCTION The growth process is a continuous process started from during conception up to adulthood, which follows a certain pattern unique to each child. Growth can be defined as increasing physical size which reflects the balance between food intake and nutritional needs of children in the process of growth (1),(2) (3). In Indonesia, shortness or stunting is a nutritional problem in infants.

The other nutritional problems in children aged 0-24 months in Indonesia were malnutrition (weight for age), severe thinness (weight for height) and severe shortness (height for age) with 5.7%, 6.8% and 19.2% respectively in 2013. Nutritional problems in children aged 0-24 months in South Sulawesi were malnutrition and severe malnutrition (weight for age), severe thinness, and thinness (height by Age) and severe shortness and shortness (height for age) which were 25%, 40% and 11% respectively (4) (5).

Triggering factors of those nutritional problems mentioned above were low quality and quantity of nutritional intake. Low macro nutrients intake were of energy and protein intakes, while for micronutrients, they were vitamin A and zinc in particular. Preventive effort to improve nutrition for individuals is through nutritional supplementation, which zinc (Zn) supplementation is one the kind and is recommended to address the growth problems in children.

From double-blind study conducted in Vietnam, by administering 10 mg Zn/ day

towards 146 children aged 4-36 months for 5 months showed that zinc supplementation increases body weight ($+ 0.5 \pm 0.1$ kg; $p < 0.05$) and body height ($+ 1.5 \pm 0.2$ Kg; $p < 0.001$) (4). Thus, in order to overcome the problem of malnutrition, particularly shortness status is through the administration of Zn supplementation.

There are several studies granting sources of zinc from natural ingredients such as fish and resulted increasing levels of Zn which was influenced by protein reserves. Provision of Zn as food can increase the levels of Zn at the time of low zinc status in children with malnutrition (1) (6). Animal source of zinc such as fish and meat can be better absorbed than those of vegetables one.

For instances, *Euthynnus affinis* contains as many as 3 mg Zn per 100gr and the *Caesionidae* contains as much as 1,5 mg Zn per 100gr, while vegetable sources are often bounded by phytate which can interfere the absorption of Zn (2),(7). Fish is a source of high quality of protein, with digestibility as much as 96%. High digestibility of protein is necessary for growth and intelligence. Fishmeal is a source of protein that has high value of amino acids.

The analysis of nutritional value of fishmeal made of *Rastrelliger kanagurta* in 100gr it consists of: protein 26.87gr, fat 18.98 g, carbohydrate 36.28g, iodine 2.67 mg, crude fiber 2.51 g and calcium 238 mg, then the fishmeal was tested for its acceptance levels among children under five. The result showed a high level of preference as a topping when eating.

For *Euthynnus affinis* in 100 g, it was obtained 30 mg of Zn, while the *Caesionidae* contained 15 mg (8). Based on the frameworks, this study aimed to analyze the provision of Zn-rich fishmeal towards levels of zinc, serum vitamin A and nutritional intake and shortness status of children aged 2-4 years after the administration of vitamin A, in Makassar South Sulawesi, Indonesia.

Methods and Materials This study used quasi experimental study. The variable included were the family's identity, the identity of the child, the nutritional status of height by age, intake of macro and micronutrients. Anthropometric measurements were done according to the standard measurement of height and weight the Ministry of Health of the Republic of Indonesia.

Macro and micro nutrient intake were obtained using a 24-hour recall with a multi-pass approach. Interviews were performed at respondents' homes independently. Respondents signed study agreements in entirety. Instruments used on the study had been previously tested at 10% of the population and was declared fit for use.

Enumerators in this study were nutritionists with diploma degree graduated from Health Polytechnic Makassar.

Enumerators were carefully trained to conduct interviews and perform anthropometric measurements. Anthropometric tools used were microtoise for height and were products of Nutritional Department of Health Polytechnic Makassar in 2010 and were in a state of good for use. Data processing and analysis of nutritional status were done using WHO Antro application, 2005.

RESULTS Macro and Micro Nutrient Intake Energy intake in group I (Vitamin A), did not increase significantly before and after the intervention ($p = 0.278$), the same occurred in group II (Vitamin A plus fishmeal) ($p = 0.237$). Significant increases in the intake of protein, fat, carbohydrates, Vitamin A and Zinc occurred in both groups ($p < 0.050$) (Table 1).

Biochemical Indicators In this study, measured biochemical indicators were serum levels of Zn, albumin, total protein and hemoglobin. Zinc serum levels increased significantly in both groups ($p < 0.050$). Albumin was not increased in group I (Vitamin A) ($p = 0.360$), but there was an increase in group II (Vitamin A plus fishmeal) ($p = 0.010$). Protein total increased in group I (Vitamin A) ($p = 0.007$), but did not in group II (Vitamin A plus fishmeal) ($p = 0.531$). Hemoglobin did not increase in group I (Vitamin A) ($p = 0.092$), but increased in group II (Vitamin A plus fishmeal) ($p = 0.009$) (Table 2)

DISCUSSION

Energy intake in group I (Vitamin A), was not increased significantly before and after the intervention ($p = 0.278$), the same occurred in group II (Vitamin A plus fishmeal) ($p = 0.237$).

Significant increases were found in the intake of protein, fat, carbohydrates, Vitamin A and Zinc and was occurred in both groups ($p < 0.050$). Energy intake is important, to support normal metabolic processes for body growth and development. Deficiency of energy intake results to growth failure. The most noticeable feature in growth failure is short stature (stunting). The entire sample in this study was children with shortness.

Food consumption history of the children with shortness begins with a low energy intake since age of infancy. To feed children with insufficient nutrients result to shortness in the long term particularly in the period of growth. Low energy intake results to low effectiveness of utilization of micronutrients.

It is important in a child's feeding intervention, to put energy intake as the primary indicator. This means that all nutritional intervention in children should be performed only when the energy intake meets nutritional needs. When energy needs do not meet

the needs, then theoretically it is difficult to obtain a positive effect on the indicator of growth and development of children.

The main difficulty in this study was to increase energy intake, while the food security in household was also poor. This study was conducted towards low-income groups.

Limitation of this study was that it did not include socio-economic variables as controlled variable. Difficulty to perform insulation in every family was the main reason.

From the results of this study, it demonstrated that the combination of vitamin A and Zinc provides better results for reducing morbidity from the diseases (9),(10),(11). In this study, measured biochemical indicators were serum levels of Zn, albumin, total protein and hemoglobin. The increase of zinc serum levels was significant in both groups ($p < 0.050$). Albumin was not increased in group I (Vitamin A) ($p = 0.360$), and there was an increase in group II (Vitamin A plus fishmeal) ($p = 0.010$). Protein total increased in group I (Vitamin A) ($p = 0.007$), but did not in group II (Vitamin A plus fishmeal) ($p = 0.531$). Hemoglobin did not increase in group I (Vitamin A) ($p = 0.092$), but did increase in group II (Vitamin A plus fishmeal) ($p = 0.009$) (Table 2) Zinc deficiency since the time of conception will affect the child's height growth.

For this reason it is important to perform zinc supplementation. The best way is through enrichment of sources of zinc. The advantage is that if the food sources of zinc are used then the sustainability of the program will be very well (12). CONCLUSION From final result of this study it can be concluded that the intervention of vitamin A supplementation can be coupled with a zinc-rich foods such as fishmeal.

ACKNOWLEDGMENTS This research can be carried out with the help of various parties; the Department of Nutrition, Health Polytechnic Makassar, and the Director that provided the facilities. The Government and the Health Department and staffs in the Municipality of Makassar and of South Sulawesi Province. It is conveyed as well to the data collectors and supervisor.

CONFLICT OF INTEREST This study did not have any conflict of interests with any party involved in the study. The study was planned, performed, monitored and the results reported scientifically and was not be intervened by another party. REFERENCE Aritonang Irianto, 2010. Menilai Status Gizi Untuk Mencapai Sehat Optimal, Leutika dengan CEBios, hlm 1-8. Barasi Maria E, 2009. At Glance Ilmu Gizi. Jakarta: Erlangga, hlm 124-129. BAPENAS,2011.

Rencana Aksi Nasional Pangan dan Gizi 2005-2015. ISBN 978-979-3767-68-9 Bentley Margaret E, dkk, 1996. Zinc Supplementation Affects The Activity Patterns of Rural

Guatemala Infants, Community and International Nutrition, American Society For Nutritional Sciences, 1333-1338. 6. Hernawati Ina, 2000, Pencegahan dan Penanggulangan Gizi Buruk, Seminar Nasional Hari Pangan Sedunia, hlm 20-30. 5. Republik Indonesia. Kementerian Kesehatan.

Laporan Hasil Riset Kesehatan Dasar Tahun 2013. Badan Penelitian dan Pengembangan Kesehatan. Jakarta, 2013. Lind Torbojorn, et al, 2003, A Community Based Randomized Controlled Trial of Iron and Zinc, Am J Clin Nutr 2003; 77: 883-890 Set Owusu Agyei, Sam Newton, Emmanuel Mahama, Lawrence Geyba Febir, Martha Ali, Kwame Adjei, Kofi Tchum, Latifa Al Hasan, Tabisile Moleah and Sherry A Tanumiharjo, Impact of Vitamin A with Zinc Supplementation on Malaria Morbidity in Ghana. Nutrition Journal 2013, 12:131 (23 September 2013).

Mustamin, dkk, 2011, Daya Terima Bubuk Gizi (Sprinkle) Pada Anak Umur 24-36 Bulan di Wilayah Kerja Puskesmas Sudiang Raya, Resbinakes Politeknik Kesehatan Kemenkes Makassar. Wapnir Raul A, 2000, Zinc Deficiency, Malnutrition and The Gastrointestinal Tract, Zinc and Health: Current Health and Future Direction, American Society For Nutritional Sciences, 2000: p 1388S – 1392S Li Chen, Yong-Fang Liu Ms/ Min Gong, Wei Jiang, Zhen Fan, et.al, 2012, Effects of vitamin A, vitamin A plus zinc, and multiple micronutrients on anemia in preschool children in Chongqing, China, Asia Pasifik Journal clinic Nutrition, Vol.21.No.1.p.3-11. Kurt Z Long, Yura Montoya, Ellen Hertzmark, Jose I Santos, and Jorge L Rosado.

A, 2011, Double-blind, Randomized, Clinical Trial of The Effect of Vitamin A and Zinc Supplementation on Diarrheal Disease and Respiratory Tract Infections in Children In Mexico City, Mexico. The American Journal Of Clinical Nutrition. Vol. 83, pp. 693-700. Ninh Nguyen Xuan, dkk, 1996, Zinc Supplementation Increases Growth and Circulating Insulin Like Growth Factor I (IGF-1) in Growth-Related Vietnamese Children, Am J Clin Nutr 1996;63:514-519 Table 1.

Macro and Micro Nutrients Intakes of Children Intake Variable _Category _Pre- Intervention _Post- Intervention _ _ _Group I (Vit. A) _Group II (Vit A + fishmeal) _Group I (Vit. A) _Group II (Vit A + fishmeal) _ _ _N _% _n _% _N _% _N _% _Energy _Good _4 _26,7 _5 _33,3 _5 _33,3 _5 _33,3 _ _Sufficient _7 _46,7 _7 _46,7 _8 _53,3 _9 _60,0 _ _Insufficient _4 _26,6 _3 _20,0

_2 _13,3 _1 _6,7 _Protein _Good _1 _6,7 _14 _93,3 _2 _13,3 _2 _13,3 _ _Sufficient _11 _73,3 _0 _0 _13 _86,7 _13 _86,7 _ _Insufficient _3 _20,0 _1 _6,7 _0 _0 _0 _0 _Fat _Good _0 _0 _0 _0 _2 _13,3 _4 _26,7 _ _Sufficient _12 _80 _11 _73,3 _13 _86,7 _9 _60,0 _ _Insufficient _3 _20 _4 _26,7 _0 _0 _2 _13,3 _Carbohydrate _Good _4 _26,7 _4 _26,7 _8

_53,8 _7 _46,7 _ _Sufficient _11 _73,3 _11 _73,3 _7 _46,7 _8 _53,8 _ _Insufficient _0 _0 _0
_0 _0 _0 _0 _0 _ _Vit.

A _Good _12 _80 _15 _100 _13 _86,7 _15 _100 _ _Sufficient _0 _0 _0 _0 _0 _0 _0 _ _
_Insufficient _3 _20 _0 _0 _2 _13,3 _0 _0 _ _Zn _Good _13 _86,7 _11 _73,3 _15 _100 _15
_100 _ _Sufficient _0 _0 _0 _0 _0 _0 _0 _ _Insufficient _2 _13,3 _4 _26,7 _0 _0 _0 _0 _ _

Table 2. Macro and Micro Nutrients Intakes Group _Intakes _Pre- Intervention, RDA
Percentage (x±SD) _Post- Intervention RDA Percentage, (x±SD) _Paired T-Test (p) _
_Vitamin A _Energy _86,73±11,98 _88,13±10,99 _0,278 _ _Protein _83,27±8,66
_88,60±6,56 _0,000 _ _Fat _81,67±6,73 _87,67±5,94 _0,000 _ _Carbohydrate
_94,13±7,16 _98,07±3,26 _0,010 _ _Vitamin A _403,33±47,91 _430,73±36,58 _0,000 _ _
_Zn _8,77±0,70 _8,95±0,71 _0,002 _ _Vitamin A Plus Fishmeal _Energy _90,13±11,62
_92,00±9,60 _0,237 _ _Protein _86,33±6,67 _90,67±5,94 _0,000 _ _Fat _84,00±8,90
_89,67±8,34 _0,000 _ _Carbohydrate _93,93±5,52 _97,4±3,31 _0,001 _ _Vitamin A
_421,67±23,88 _446,67±6,99 _0,003 _ _Zn _8,60±0,68 _8,91±0,51 _0,020 _ _Total
_Energy _88,74±11,72 _90,00±10,32 _0,100 _ _Protein _84,0±7,75 _89,63±6,23 _0,000 _
_ _Fat _82,83±7,84 _88,66±7,18 _0,000 _ _Carbohydrate _94,03±6,28 _97,73±3,24 _0,000
_ _Vitamin A _412,5±38,34 _438,70±27,11 _0,000 _ _Zn _8,68±0,68 _8,92±0,61 _0,001 _
_ Table 3.

Biochemical Indicators Pre and Post Intervention Vitamin A _Zn _53,40±8,20 _63,46±2,29
_0,000 _ _Albumin _5,53±8,18 _3,58±0,82 _0,360 _ _Total Protein _3,28±0,70
_3,88±0,91 _0,007 _ _Hemoglobin _10,72±2,08 _11,80±0,77 _0,092 _ _Vitamin A Plus
Fishmeal _Zn _53,73±10,73 _63,73±10,20 _0,016 _ _Albumin _3,22±0,63 _3,90±0,47
_0,010 _ _Total Protein _3,70±1,37 _4,02±0,58 _0,531 _ _Hemoglobin _10,13±1,47
_11,73±1,38 _0,009 _ _Total _Zn _53,56±9,39 _63,60±7,36 _0,000 _ _Albumin _4,38±5,82
_3,74±0,68 _0,547 _ _Total Protein _3,53±1,1 _3,95±0,75 _0,056 _ _Hemoglobin
_10,43±1,77 _11,76±1,10 _0,002 _ _

INTERNET SOURCES:

2% -

[http://gssrr.org/index.php?journal=JournalOfBasicAndApplied&page=article&op=view
&path%5B%5D=4999](http://gssrr.org/index.php?journal=JournalOfBasicAndApplied&page=article&op=view&path%5B%5D=4999)

<1% - <http://slimdoggy.com/dog-food-protein-part-1/>

<1% - <http://fkg.jtam.unlam.ac.id/index.php/dentin/article/view/169>

<1% - http://gadi.agric.za/Agric/Vol14No1_2014/semen.php

<1% - https://www.researchgate.net/profile/Sonia_Philippi

1% - <https://link.springer.com/article/10.1007%2Fs12098-014-1483-7>

<1% - https://en.wikipedia.org/wiki/Child_development

<1% - https://en.wikipedia.org/wiki/Basal_metabolic_rate
<1% -
<https://www.omicsonline.org/open-access/nutritional-intake-biochemical-status-and-anthropometric-values-of-stunting-children-after-added-zinc-on-vitamin-a-supplementation-2168-9652-S5-008.pdf>
<1% -
<https://www.epi.org/publication/broad-based-wage-growth-is-a-key-tool-in-the-fight-against-poverty/>
1% -
<https://www.omicsonline.org/open-access/obstacles-of-breastfeeding-contributed-to-stunted-children-status-in-barru-regency-south-sulawesi-2168-9652-S5-004.pdf>
1% -
<http://gssrr.org/index.php?journal=JournalOfBasicAndApplied&page=article&op=download&path%5B%5D=6177&path%5B%5D=2998>
<1% - <http://www.mdpi.com/2072-6643/8/7/428/htm>
<1% - <https://academic.oup.com/jcem/article/82/5/1549/2823430>
<1% - <http://europepmc.org/articles/PMC3060650/reload=2>
<1% - <https://quizlet.com/31741222/nutrition-chapter-9-minerals-flash-cards/>
<1% - <https://www.feedipedia.org/node/72>
<1% - https://www.researchgate.net/profile/Sara_Wilcox
<1% - <https://www.tandfonline.com/doi/full/10.1080/00220388.2016.1228881>
<1% - https://www.researchgate.net/profile/Allen_Knehans
<1% - <https://jasbsci.biomedcentral.com/articles/10.1186/s40104-017-0194-0>
<1% - <https://www.onlinelibrary.wiley.com/doi/full/10.1002/jcsm.12237>
<1% - https://www.researchgate.net/profile/Hilde_Brekke
<1% -
<https://www.encyclopedia.com/science-and-technology/biochemistry/biochemistry/growth-hormone>
<1% -
<https://journals.lww.com/acsm-msse/Pages/ArticleViewer.aspx?year=2009&issue=03000&article=00027&type=Fulltext>
<1% - <http://www.gpo.gov/fdsys/pkg/FR-2017-10-20/xml/FR-2017-10-20.xml>
<1% - <http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-2909.133.2.205>
<1% - <http://www.mdpi.com/2072-6643/9/6/624/htm>
<1% - <https://www.scribd.com/document/234938958/who-ehc-221-1>
<1% - <https://www.scribd.com/document/333217424/4c-Full-Proceeding>
<1% - <http://en.amda.or.jp/articlelist/?tags=66>
<1% -
<http://indianexpress.com/article/opinion/columns/no-proof-required-conflict-of-interest-anyone/>

1% -

https://www.researchgate.net/profile/Elena_Hurtado3/publication/14017440_Zinc_Supplementation_Affects_the_Activity_Patterns_of_Rural_Guatemalan_Infants/links/571fb4b308aead26e71b704a/Zinc-Supplementation-Affects-the-Activity-Patterns-of-Rural-Guatemalan-Infants.pdf

1% -

<http://biofarmaka.ipb.ac.id/biofarmaka/2014/Riskedas2007%20-%20Province%20Report%2016%20SUMSEL.pdf>

1% - <https://www.sciencedirect.com/science/article/pii/S0899900710001280>

<1% - <https://www.hindawi.com/journals/ipid/2010/753512/ref/>

1% - <http://journals.sagepub.com/doi/abs/10.1177/0884533615570376>

<1% -

http://journals.lww.com/acsm-msse/Fulltext/2014/05001/C_34_Free_Communication_Poster__Sports_Nutrition,.85.aspx